

## LIST OF APPENDICES

	<b>Page</b>	
Appendix 3.2.2.1	Field strains of <i>Aedes aegypti</i> and <i>Aedes albopictus</i> were obtained from ovitrap collection meanwhile <i>Culex quinquefasciatus</i> field strain was collected by bare-leg catch method from Faculty of Science, University Malaya	262
Appendix 3.4.1	Preparation for amount of insecticide at different concentration	269
Appendix 3.5.1	Various concentration (mg/L) of insecticide for larval bioassay to determine LC50 from percentage of mortality	270
Appendix 3.7.1	Probit analysis print out for larval bioassay	271
Appendix 3.8.1.1	LC <sub>50</sub> (mg/L) values in insecticide test of early fourth instar larvae of laboratory and resistant strains of <i>Culex quinquefasciatus</i> of 10 subsequent generation exposed continuously for 24 hours to malathion	272
Appendix 3.8.1.2	LC <sub>50</sub> (mg/L) values in insecticide test of early fourth instar larvae of laboratory and resistant strains of <i>Culex quinquefasciatus</i> of 10 subsequent generation exposed continuously for 24 hours to permethrin	273
Appendix 3.8.2.1	LC <sub>50</sub> (mg/L) values in insecticide test of early fourth instar larvae of laboratory and resistant strains of <i>Aedes aegypti</i> of 10 subsequent generation exposed continuously for 24 hours to malathion	274
Appendix 3.8.2.2	LC <sub>50</sub> (mg/L) values in insecticide test of early fourth instar larvae of laboratory and resistant strains of <i>Aedes aegypti</i> of 10 subsequent generation exposed continuously for 24 hours to permethrin	275
Appendix 3.8.2.3	LC <sub>50</sub> (mg/L) values in insecticide test of early fourth instar larvae of laboratory and resistant strains of <i>Aedes aegypti</i> of 10 subsequent generation exposed continuously for 24 hours to temephos	276
Appendix 3.9.1	List of diagnostic dosage (mg/L) recommended by WHO 1992	277
Appendix 4.3.1	Probit analysis print out for adult bioassay	278

Appendix 4.3.2	Adult bioassay form to record adult mortality	278
Appendix 5.3.1	Print out of the OD readings for <i>Cx. quinquefasciatus</i> malathion strain	279
Appendix 5.3.2	Print out of the OD readings for <i>Cx. quinquefasciatus</i> permethrin strain	280
Appendix 5.3.3	Print out of the OD readings for <i>Ae. albopictus</i> temephos strain	281
Appendix 5.3.4	Print out of one way analysis of variance (ANOVA) to compare the enzyme expression levels between life stages of insecticide selected strain to susceptible strain	282
Appendix 5.3.5	Represent print out of the analysis of correlation between mean enzyme activity and LT <sub>50</sub> values using spearman rank-order correlation	283
Appendix 5.4.1.1	Optical density (OD) of non-specific esterases towards life stages of <i>Culex quinquefasciatus</i> susceptible control and malathion selected ten subsequent test populations against $\alpha$ – naphthyl acetate	285
Appendix 5.4.1.2	Optical density (OD) of non-specific esterases towards life stages of <i>Culex quinquefasciatus</i> susceptible control and permethrin selected ten subsequent test populations against $\alpha$ – naphthyl acetate	288
Appendix 5.4.2.1	Optical density (OD) of non-specific esterases towards life stages of <i>Aedes aegypti</i> susceptible control and malathion selected ten subsequent test populations against $\alpha$ – naphthyl acetate	291
Appendix 5.4.2.2	Optical density (OD) of non-specific esterases towards life stages of <i>Aedes aegypti</i> susceptible control and permethrin selected ten subsequent test populations against $\alpha$ – naphthyl acetate	294
Appendix 5.4.2.3	Optical density (OD) of non-specific esterases towards life stages of <i>Aedes aegypti</i> susceptible control and temephos selected ten subsequent test populations against $\alpha$ – naphthyl acetate	297

Appendix 5.4.3.1	Optical density (OD) of non-specific esterases towards life stages of <i>Aedes albopictus</i> susceptible control and malathion selected ten subsequent test populations against $\alpha$ – naphthyl acetate	300
Appendix 5.4.3.2	Optical density (OD) of non-specific esterases towards life stages of <i>Aedes albopictus</i> susceptible control and permethrin selected ten subsequent test populations against $\alpha$ – naphthyl acetate	302
Appendix 5.4.3.3	Optical density (OD) of non-specific esterases towards life stages of <i>Aedes albopictus</i> susceptible control and temephos selected ten subsequent test populations against $\alpha$ – naphthyl acetate	306
Appendix 5.5.1.1	Percentage of non-specific esterase activity in life stages of field strains against $\alpha$ – naphthyl acetate	309
Appendix 5.5.1.2	Percentage of non-specific esterase activity in life stages of <i>Culex quinquefasciatus</i> susceptible control and malathion selected ten subsequent test populations against $\alpha$ – naphthyl acetate	312
Appendix 5.5.1.3	Percentage of non-specific esterase activity in life stages of <i>Culex quinquefasciatus</i> susceptible control and permethrin selected ten subsequent test populations against $\alpha$ – naphthyl acetate	315
Appendix 5.5.1.4	Percentage of non-specific esterase activity in life stages of <i>Aedes aegypti</i> susceptible control and malathion selected ten subsequent test populations against $\alpha$ – naphthyl acetate	318
Appendix 5.5.1.5	Percentage of non-specific esterase activity in life stages of <i>Aedes aegypti</i> susceptible control and permethrin selected ten subsequent test populations against $\alpha$ – naphthyl acetate	321
Appendix 5.5.1.6	Percentage of non-specific esterase activity in life stages of <i>Aedes aegypti</i> susceptible control and temephos selected ten subsequent test populations against $\alpha$ – naphthyl acetate	324

Appendix 5.5.1.7	Percentage of non-specific esterase activity in life stages of <i>Aedes albopictus</i> susceptible control and malathion selected ten subsequent test populations against $\alpha$ – naphthyl acetate	327
Appendix 5.5.1.8	Percentage of non-specific esterase activity in life stages of <i>Aedes albopictus</i> susceptible control and permethrin selected ten subsequent test populations against $\alpha$ – naphthyl acetate	330
Appendix 5.5.1.9	Percentage of non-specific esterase activity in life stages of <i>Aedes albopictus</i> susceptible control and temephos selected ten subsequent test populations against $\alpha$ – naphthyl acetate	333