

LIST OF TABLES

		Page
Table 1.1	Factors influencing development of resistance to insecticides in a population	5
Table 3.1	Resistant (selection pressure- treated with insecticides) strains of mosquitoes which were treated with diagnostic concentration of stock solution A and stock solution B. Preparation of stock solution of insecticides shown in Table (3.4.2)	42
Table 3.2	Preparation of diagnostic concentrations (insecticides) of stock solution A and stock solution B	42
Table 3.3	LC ₅₀ (mg/L) values in insecticide test of early fourth instar larvae of laboratory and resistant strains of <i>Aedes albopictus</i> of 10 subsequent generation exposed continuously for 24 hours to malathion	53
Table 3.4	LC ₅₀ (mg/L) values in insecticide test of early fourth instar larvae of laboratory and resistant strains of <i>Aedes albopictus</i> of 10 subsequent generation exposed continuously for 24 hours to permethrin	54
Table 3.5	LC ₅₀ (mg/L) values in insecticide test of early fourth instar larvae of laboratory and resistant strains of <i>Aedes albopictus</i> of 10 subsequent generation exposed continuously for 24 hours to temephos	55
Table 3.6	Summary of mean \pm S.E. for LC 50 of selected strains obtained for malathion, permethrin and temephos on fourth instar larvae	56
Table 4.1	WHO standard exposure time to the diagnostic concentration of insecticide impregnated filter paper	68
Table 4.2	LT ₅₀ (min) values and 24 hours post-exposure mortality of malathion resistant <i>Culex quinquefasciatus</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of malathion 5.0%	78
Table 4.3	Comparative mean percentage for 24 hrs mortality for ten generations of selection pressure for malathion resistant <i>Culex quinquefasciatus</i> treated with diagnostic dosage 5.0% malathion	79

Table 4.3	Comparative mean percentage for 24 hrs mortality of selection pressure for malathion field strain <i>Culex quinquefasciatus</i> treated with diagnostic dosage 5.0% malathion	79
Table 4.4	LT ₅₀ (min) values and 24 hours post-exposure mortality of permethrin resistant <i>Culex quinquefasciatus</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of permethrin 0.75%	81
Table 4.5	Comparative mean percentage for 24 hrs mortality for ten generations of selection pressure for permethrin resistant <i>Culex quinquefasciatus</i> treated with diagnostic dosage 0.75% permethrin	82
Table 4.6	Comparative mean percentage for 24 hrs mortality of selection pressure for permethrin field strain <i>Culex quinquefasciatus</i> treated with diagnostic dosage 0.75% permethrin	82
Table 4.7	Cross-resistance susceptibility LT ₅₀ (min) values and 24 hours post-exposure mortality of malathion resistant <i>Culex quinquefasciatus</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of propoxur 0.1%	84
Table 4.8	Comparative mean percentage for 24 hrs mortality for ten generations of selection pressure for malathion resistant <i>Culex quinquefasciatus</i> treated for cross-resistance with diagnostic dosage 0.1 % propoxur	85
Table 4.9	Comparative mean percentage for 24 hrs mortality of selection pressure for field strain <i>Culex quinquefasciatus</i> treated for cross-resistance with diagnostic dosage 0.1 % propoxur	85
Table 4.10	Cross-resistance susceptibility LT ₅₀ (min) values and 24 hours post-exposure mortality of permethrin resistant <i>Culex quinquefasciatus</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of propoxur 0.1%	87
Table 4.11	Comparative mean percentage for 24 hrs mortality for ten generations of selection pressure for permethrin resistant <i>Culex quinquefasciatus</i> treated for cross-resistance with diagnostic dosage 0.1 % propoxur	88

Table 4.12	LT ₅₀ (min) values and 24 hours post-exposure mortality of malathion resistant <i>Aedes aegypti</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of malathion 5.0%	95
Table 4.13	Comparative mean percentage for 24 hrs mortality for ten generations of selection pressure for malathion resistant <i>Aedes aegypti</i> with diagnostic dosage 5.0% malathion	96
Table 4.14	Comparative mean percentage for 24 hrs mortality of selection pressure for malathion field strain <i>Aedes aegypti</i> treated with diagnostic dosage 5.0% malathion	96
Table 4.15	LT ₅₀ (min) values and 24 hours post-exposure mortality of permethrin resistant <i>Aedes aegypti</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of permethrin 0.75%	98
Table 4.16	Comparative mean percentage for 24 hrs mortality for ten generations of selection pressure for permethrin resistant <i>Aedes aegypti</i> with diagnostic dosage 0.75% permethrin	99
Table 4.17	Comparative mean percentage for 24 hrs mortality of selection pressure for permethrin field strain <i>Aedes aegypti</i> treated with diagnostic dosage 0.75% permethrin	99
Table 4.18	Cross-resistance susceptibility LT ₅₀ (min) values and 24 hours post-exposure mortality of malathion resistant <i>Aedes aegypti</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of propoxur 0.1%	101
Table 4.19	Comparative mean percentage for 24 hrs mortality for ten generations of selection pressure for malathion resistant <i>Aedes aegypti</i> treated for cross-resistance with diagnostic dosage 0.1 % propoxur	102
Table 4.20	Comparative mean percentage for 24 hrs mortality of selection pressure for field strain <i>Aedes aegypti</i> treated for cross-resistance with diagnostic dosage 0.1 % propoxur	102
Table 4.21	Cross-resistance susceptibility LT ₅₀ (min) values and 24 hours post-exposure mortality of permethrin resistant <i>Aedes aegypti</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of propoxur 0.1%	104

Table 4.22	Comparative mean percentage for 24 hrs mortality for ten generations of selection pressure for permethrin resistant <i>Aedes aegypti</i> treated for cross-resistance with diagnostic dosage 0.1 % propoxur	105
Table 4.23	Cross-resistance susceptibility LT ₅₀ (min) values and 24 hours post-exposure mortality of temephos resistant <i>Aedes aegypti</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of propoxur 0.1%	107
Table 4.24	Comparative mean percentage for 24 hrs mortality for ten generations of selection pressure for temephos resistant <i>Aedes aegypti</i> treated for cross-resistance with diagnostic dosage 0.1 % propoxur	108
Table 4.25	LT ₅₀ (min) values and 24 hours post-exposure mortality of malathion resistant <i>Aedes albopictus</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of malathion 5.0%	115
Table 4.26	Comparative mean percentage for 24 hrs mortality for ten generations of selection pressure for malathion resistant <i>Aedes albopictus</i> treated with diagnostic dosage 5.0% malathion	116
Table 4.27	Comparative mean percentage for 24 hrs mortality of selection pressure for malathion field strain <i>Aedes albopictus</i> treated with diagnostic dosage 5.0% malathion	116
Table 4.28	LT ₅₀ (min) values and 24 hours post-exposure mortality of permethrin resistant <i>Aedes albopictus</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of permethrin 0.75%	118
Table 4.29	Comparative mean percentage for 24 hrs mortality for ten generations of selection pressure for permethrin resistant <i>Aedes albopictus</i> treated with diagnostic dosage 0.75 % permethrin	119
Table 4.30	Comparative mean percentage for 24 hrs mortality of selection pressure for permethrin field strain <i>Aedes albopictus</i> treated with diagnostic dosage 0.75% permethrin	119

Table 4.31	Cross-resistance susceptibility LT ₅₀ (min) values and 24 hours post-exposure mortality of malathion resistant <i>Aedes albopictus</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of propoxur 0.1%	121
Table 4.32	Comparative mean percentage 24 hrs mortality for ten generations of selection pressure for malathion resistant <i>Aedes albopictus</i> treated for cross-resistance with diagnostic dosage 0.1 % propoxur	122
Table 4.33	Comparative mean percentage 24 hrs mortality for ten generations of selection pressure for field strain <i>Aedes albopictus</i> treated for cross-resistance with diagnostic dosage 0.1 % propoxur	122
Table 4.34	Cross-resistance susceptibility LT ₅₀ (min) values and 24 hours post-exposure mortality of permethrin resistant <i>Aedes albopictus</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of propoxur 0.1%	124
Table 4.35	Comparative mean percentage for 24 hrs mortality for ten generations of selection pressure for permethrin resistant <i>Aedes albopictus</i> treated for cross-resistance with diagnostic dosage 0.1 % propoxur	125
Table 4.36	Cross-resistance susceptibility LT ₅₀ (min) values and 24 hours post-exposure mortality of temephos resistant <i>Aedes albopictus</i> adult female mosquitoes of 10 subsequent generation exposed against WHO diagnostic dosage of propoxur 0.1%	127
Table 4.37	Comparative mean percentage for 24 hrs mortality for ten generations of selection pressure for temephos resistant <i>Aedes albopictus</i> treated for cross-resistance with diagnostic dosage 0.1 % propoxur	128
Table 6.1	Summary of non-specific esterase activity bands in <i>Cx. quinquefasciatus</i> malathion resistant strain	197
Table 6.2	Summary of non-specific esterase activity bands in <i>Cx. quinquefasciatus</i> permethrin resistant strain	197
Table 6.3	Summary of non-specific esterase activity bands in <i>Cx. quinquefasciatus</i> susceptible strain	198

Table 6.4	Summary of non-specific esterase activity bands in <i>Cx. quinquefasciatus</i> field strain	198
Table 6.5	Summary of non-specific esterase activity bands in <i>Ae. aegypti</i> malathion resistant strain	204
Table 6.6	Summary of non-specific esterase activity bands in <i>Ae. aegypti</i> permethrin resistant strain	204
Table 6.7	Summary of non-specific esterase activity bands in <i>Ae. aegypti</i> temephos resistant strain	205
Table 6.8	Summary of non-specific esterase activity bands in <i>Ae. aegypti</i> susceptible strain	205
Table 6.9	Summary of non-specific esterase activity bands in <i>Ae. aegypti</i> field strain	206
Table 6.10	Summary of non-specific esterase activity bands in <i>Ae. albopictus</i> malathion resistant strain	213
Table 6.11	Summary of non-specific esterase activity bands in <i>Ae. albopictus</i> permethrin resistant strain	213
Table 6.12	Summary of non-specific esterase activity bands in <i>Ae. albopictus</i> temephos resistant strain	214
Table 6.13	Summary of non-specific esterase activity bands in <i>Ae. albopictus</i> susceptible strain	214
Table 6.14	Summary of non-specific esterase activity bands in <i>Ae. albopictus</i> field strain	215